

Updated COVID-19 Quick Reference Public Health Guidance on Testing and Clearance

Update to York Region health care professionals, LTCH/RH and community living facilities as of May 14, 2020

This information can be used to help guide decision making on testing and clearance of contacts of cases or individuals suspected or confirmed to have COVID-19.

Approaches to Clearing Cases

For each scenario, isolation after symptom onset should be for the duration specified provided that the individual is afebrile, and symptoms are improving. Absence of cough is not required for those known to have chronic cough or who are experiencing reactive airways post-infection. Once a case is discharged from isolation, their case status should be updated to 'resolved'. If an individual has tested positive but has never had symptoms, isolation recommendations should be based on date of specimen collection. After an individual completes their isolation period, they should continue to practice physical distancing measures as recommended for everyone at this time. The guidance below is based on the observation that some people with more severe illness may have prolonged detection of viral RNA which may indicate the potential for longer viral shedding; for ease of use, "severe illness" has been defined as having required hospitalization for their COVID-19 illness.

	When to Use	Instructions
<p>Non-Test Based Approach</p> <p>Waiting 14 days from symptom onset (or 14 days from when swab was taken if persistently asymptomatic)</p>	<p>Appropriate for most individuals who have recovered from mild to moderate illness (i.e., never hospitalized) including:</p> <ul style="list-style-type: none"> • Health care workers (unless otherwise directed by their employer/Occupational Health and Safety) • individuals who live in congregate settings (e.g., long-term care homes, shelters) 	<p>Can discontinue isolation at 14 days after symptom onset (or 14 days from positive test collection date if never had symptoms), provided that the individual is afebrile and symptoms are improving. Absence of cough is not required for those known to have chronic cough or who are experiencing reactive airways post-infection.</p>
<p>Test Based Approach</p> <p>Two consecutive negative specimens collected at least 24 hours apart.</p>	<p>To remove individuals from isolation who had severe illness (specifically, were hospitalized for their COVID-19 illness), and</p> <ul style="list-style-type: none"> - who remain in hospital after symptom improvement; OR - who are being discharged from hospital to continue isolating in a congregate living setting (e.g., long-term care homes, shelters) 	<p>Continue isolation until 2 consecutive negative specimens collected at least 24 hours apart.</p> <ul style="list-style-type: none"> - Clearance testing may begin after the individual has become afebrile and symptoms are improving. - Absence of cough is not required for those known to have chronic cough or who are experiencing reactive airways post-infection. - If swab remains positive, test again in approximately 3-4 days. If swab is negative, re-test in 1-2 days (and at least 24 hours apart). - Tick the box labelled 'For clearance of disease' on the PHO Laboratory COVID-19 Test Requisition, or clearly write this on the requisition if submitting to another laboratory

Source: http://www.health.gov.on.ca/en/pro/programs/publichealth/coronavirus/docs/2019_testing_clearing_cases_guidance.pdf

Notes: If test based clearance is not feasible in any scenario, the non-test based clearance approach may be used. Individuals who were hospitalized and are being discharged home, can be cleared from isolation using a non-test based approach.

Recommendations for Health Care Workers (HCW) Return to Work

- HCWs should follow isolation and clearance with a non-test based approach unless they have required hospitalization during the course of their illness, in which case a test based approach is preferred.



Some HCWs may be directed to have test based clearance by their employer/Occupational Health and Safety.

- Symptomatic HCWs awaiting testing results must be off work.
- Asymptomatic HCWs awaiting testing results may continue to work using the appropriate precautions recommended by the facility, which will depend on the reason for testing.

In **exceptional circumstances** a HCW may be allowed to return to work before the above guidelines would suggest where clinical care would be severely compromised without additional staffing. This return to work may be considered under work self-isolation recognizing the staff may still be infectious.

Work self-isolation means maintaining self-isolation measures outside of work for 14 days from symptom onset (or 14 days from positive specimen collection date if consistently asymptomatic) to avoid transmitting to household members or other community contacts. While at work, the HCW should adhere to universal masking recommendations, maintain physical distancing (remaining greater than 2 metres (6 feet) from others) except when providing direct care, and performing meticulous hand hygiene. These measures at work are required to continue until non-test based clearance (or test based clearance if required by employer/Occupational Health and Safety). The HCW should ideally be cohorted to provide care for COVID-19 positive patients/residents if possible. The HCW on work self-isolation should not work in multiple locations.

Symptoms	Test Result	Instructions
Yes	Positive	<ul style="list-style-type: none"> • Work self-isolation could start after a minimum of 72 hours after illness resolving, defined as resolution of fever and improvement in respiratory and other symptoms
Yes	Negative	<ul style="list-style-type: none"> • May return to work 24 hours after symptom resolution • If the HCW was self-isolating due to an exposure at the time of testing, return to work should be under work self-isolation until 14 days from last exposure .
Never symptomatic at time of test	Positive	<ul style="list-style-type: none"> • If there has been a recent potential exposure (e.g., tested as part of an outbreak investigation or other close contact to a case), work self-isolation (i.e., return to work) could start after a minimum of 72 hours from the positive specimen collection date to ensure symptoms have not developed in that time, as the positive result may represent early identification of virus in the pre-symptomatic period • If there has been no known recent potential exposures (e.g., tested as part of surveillance and no other cases detected in the facility or on the unit/floor, depending on the facility size), there is no minimum time off from the positive specimen collection date as it is unclear when in the course of illness the positive result represents (i.e., consistently asymptomatic HCWs can continue working in work self-isolation until 14 days from specimen collection date).

Additional Notes on Clearance

Follow-up screening or testing is **not required** for clearance of individuals with mild or moderate COVID-19 illness. COVID-19 testing can be sensitive, and has, at times, identified remaining viral RNA when an individual is no longer infectious. York Region Public Health does not facilitate retesting; any retesting would have to be supported by the individual’s attending physician.

Questions?

Call our dedicated health professional COVID-19 line at **1-877-464-9675 ext. 77280** (8:30 a.m. to 8 p.m., seven days a week, after hours call 905-953-6478). Continue to visit york.ca/healthprofessionals, york.ca/covid19 and Ontario.ca/covid19 for up to date information on COVID-19.