


TWO STEPS TO HELP
SAVE A LIFE
OPIOID OVERDOSE
DURING COVID-19

STEP 1 :
Look for signs of an **OVERDOSE**




Not moving and
can't be woken




Slow or not
breathing




Blue or purple lips
and nails



Tiny pupils



Cold or clammy skin



Choking, gurgling
sounds or snoring

Substance Use Prevention and Harm Reduction

Visit: york.ca/opioids or

Call: 1-877-464-9675 ext. 76683

TTY: 1-866-512-6228

Email: substance@york.ca

STEP 2:

Follow these steps when giving NALOXONE



Tap and shout



Call 911




Give naloxone: 1 nasal spray*
or injection into arm or leg



Chest compressions**



Is it working?
YES



NO improvement? Give naloxone
again, continue chest compressions
and check in 2 - 3 minutes



**PUT PERSON IN
RECOVERY POSITION (LEFT SIDE)**

IF: they begin breathing on their own or
if you have to leave them alone.

***GIVING NASAL NALOXONE DURING COVID-19 DOES NOT PRODUCE AEROSOLS.
**AT THIS TIME WE DO NOT RECOMMEND GIVING RESCUE BREATHS. PERFORMING CHEST
COMPRESSIONS DURING CPR IS NOT DANGEROUS.**

FOR MORE INFORMATION:

York Region Public Health Substance Use Prevention and Harm Reduction program:

substance@york.ca or 1-877-464-9675 ext. 76683

Report Bad Drugs: york.ca/reportbaddrugs

Connex Ontario: connexontario.ca or 1-866-531-2600